

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS
BEFORE OPENING
OR USING



WEEDFORCE®

Gusto®

WOODY & NOXIOUS
WEED HERBICIDE

**ACTIVE CONSTITUENTS: 300 g/L TRICLOPYR PRESENT AS THE BUTOXYETHYL
ESTER 100 g/L PICLORAM PRESENT AS HEXYLOXYPROPYLAMINE SALT**

GROUP 4 HERBICIDE

For control of a range of environmental and noxious woody and herbaceous
weeds as specified in the Directions for Use table.

**IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT.
SHAKE WELL BEFORE USE**

WEEDFORCE Pty Ltd ACN 602 207 152 
2072 Logan Rd Mt Gravatt QLD 4122
P: 0488 997 923 www.weedforce.com.au

APVMA Approval No:
83353/111562

© Registered Trademark of Weed Force Pty Ltd © Weed Force Pty Ltd 2017

WEED FORCE GUSTO WOODY & NOXIOUS WEED HERBICIDE

DIRECTIONS FOR USE

INDEX

RESTRAINTS.....	4
SECTION 1- WOODY WEED & NOXIOUS WEED SITUATIONS	
TABLE A: HIGH VOLUME SPRAYING (HAND GUN)	5
TABLE B: AERIAL APPLICATION	12
TABLE C: CONTROLLED DROPLET APPLICATION (CDA)	14
TABLE D: LOW VOLUME HIGH CONCENTRATE APPLICATION TECHNIQUES (GAS POWERED GUN, SPRINKLER SPRAYER)	14
TABLE E: BOOM APPLICATION	16
SECTION 2 - FALLOW SITUATIONS	
TABLE A: BOOM APPLICATION	16
TABLE B: BLANKET WIPER APPLICATION	18
GENERAL INSTRUCTIONS	
MINIMUM RECROPPING PERIODS	19
COMPATIBILITY	20
MIXING	20
APPLICATION & EQUIPMENT	
SECTION 1: WOODY WEED SITUATIONS	20
SECTION 2: FALLOW SITUATIONS	21
CLEANING SPRAY EQUIPMENT	22
RESISTANT WEEDS WARNING.....	22
PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.....	22
PROTECTION OF LIVESTOCK	23
PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT	23
STORAGE AND DISPOSAL.....	23
SMALL SPILL MANAGEMENT	23
SAFETY DIRECTIONS	23
FIRST AID	24
SAFETY DATA SHEET (SDS)	24
CONDITIONS OF SALE	24

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.

DO NOT spray if rain is likely within one hour or if foliage is wet from rain or dew.

DO NOT burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.

DO NOT apply by aerial application in wind in excess of 15km/hr and air temperatures above 35°C.

In areas prone to flooding treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.

1 WOODY WEED SITUATIONS

Table A: High Volume Spraying

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ 100L WATER	CRITICAL COMMENTS
African boxthorn	Less than 2m tall	All States	500mL	Apply when bushes have good leaf cover, growth and no leaf fall
<i>Angophora</i> spp.	1 to 3m tall		350mL	
Australian blackthorn	Less than 2m tall		500mL	Apply from late spring to early autumn.
<i>Banksia</i> spp.	1 to 3m tall		350mL	
Biddy bush (Chinese shrub) (Sifton bush)	Autumn when actively growing	ACT and NSW only	500mL	Add a 100% concentrate non-ionic surfactant at 125mL/100L of water for best results.
Blackberry in association with: Docks Ragwort Smartweed Thistles	Late spring to autumn	All States except NT	350mL OR 500mL	Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry. Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before respraying with Gusto Woody & Noxious Weed Herbicide.
Blue heliotrope	Flowering	NSW and Qld only	500mL	Apply in a minimum spray volume of 1250L/ha.
Brooms: Cape English Flax leaf Montpellier	Spring to mid-summer prior to pod formation	All States except NT	250mL	Apply as thorough foliage spray.
	Autumn to winter		350mL	
Camphor laurel	Less than 2m tall	All States		
	Above 2m tall		500mL	
<i>Casuarina</i> spp.	1 to 3m tall		350mL	

Table A: High Volume Spraying

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY			AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY	
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ 100L WATER	CRITICAL COMMENTS
Chinese apple	Less than 2m tall	Qld & WA only	350mL	Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Cockspur thorn	Spring to autumn	ACT, NSW & Qld only		Apply as a thorough foliage spray.
Common sensitive plant	Any time when actively growing	NT, Qld & WA only	200mL	To avoid leaves closing during application, spray plants while moving forward. Add a 100% concentrate non-ionic surfactant (at 100mL/100L of water) for best results.
Crofton weed	Spring to autumn	ACT, NSW & Qld only	350mL	Apply as a thorough foliage spray.
Eastern cotton bush (<i>Maireana microphylla</i>)	Spring to autumn	NSW & Qld only	500mL	Add Uptake Spraying Oil at 500mL/100L of water. Some bushes may require a follow-up spray to control regrowth.
<i>Eucalyptus</i> spp.	Seedling and regrowth from small lignotubers, 1 to 3m tall	All States	350mL OR 500mL	Apply the high rate where difficult to control species of <i>Eucalyptus</i> regrowth is present. Addition of an adjuvant may improve results – contact Weed Force for details.
Galenia	Fresh growth in spring to summer	NSW only	500mL	Use 2000L of spray mixture/ha. Penetration of thick clumps may be difficult and respraying may be necessary. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Giant bramble	Spring to autumn	NT, Qld & WA only		
Gorse	1 to 1.5m tall	All States except NT	250mL	Spring and Summer treatment only. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
	Over 1.5m tall or autumn treatment		350mL	Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
	Winter treatment		500mL	Brownout may not be complete until summer. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Groundsel bush (<i>Baccharis halimifolia</i>)	1 to 1.5m tall in spring and summer	All States except NT	250mL	Apply as a thorough foliar spray.
	Over 1.5m tall or autumn treatment		350mL	

Table A: High Volume Spraying
See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY			AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY	
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ 100L WATER	CRITICAL COMMENTS
Green cestrum	Late spring to early autumn	ACT, NSW and Qld only	500mL	One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high. Apply from late spring to early autumn.
Hawthorn	Less than 2m tall	All States	350mL	Apply as a thorough foliar spray. Add one of the following adjuvants when using 350mL rate: Uptake Spraying Oil @ 0.5% v/v. Pulse Penetrant
Horehound	Pre-flowering	NSW and Qld only		
Japanese sunflower	Up to 1m tall in summer to autumn 1 to 2m tall in summer to autumn		All States	500 or 750mL
Lantana (<i>Lantana camara</i>) (<i>Lantana montevidensis</i>)		Pre-flowering		Qld only
Lion Tail (<i>Leonotis nepetifolia</i>)	Any time of year with good leaf cover and soil moisture	NT and Qld only	350mL	Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results – contact Weed Force for details.
Limebush	At flowering	Vic only	500mL	For optimum results, add Pulse Penetrant at 200mL/100L of spray. Thoroughly wet foliage, stems and soil around the base of the plants.
Mesquite (<i>Prosopis</i> spp.) <i>Prosopis velutina</i>	Seedling, full leaf and flowering before podding	NSW, NT, Qld and WA only	350mL	DO NOT spray plants bearing pods. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results. Apply as a thorough foliar spray.
		Qld only	670mL	
Mistflower	Spring to autumn	ACT, NSW and Qld only	500mL	Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results. Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.
Mother-of-millions	Flowering	NSW and Qld only		
Paddy's lucerne	Active growth	NSW only	350mL	Add Uptake Spray Oil at 500mL/100L water. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.
Parkinsonia	Under 2m tall	NT, Qld and WA only		
Prickly pear (common) Smooth tree pear	Active phyllode growth	All States	500mL	Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.

Table A: High Volume Spraying
See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY			AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY	
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ 100L WATER	CRITICAL COMMENTS
Rubber vine (Not infected with rust)	Up to 1.5m tall at flowering	NT, Qld and WA only	350mL	Spray all leaves and stems just to the point of runoff and thoroughly spray the base of the plant. With larger, more dense stands regrowth may occur. Subsequent control of any regrowth should be done by basal bark spraying.
	Dense stands greater than 1.5m tall at flowering		500mL	
Siam weed	Active growth	Qld and WA only	350mL	Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Sicklepod	Up to flowering	NT, Qld and WA only	200mL	DO NOT apply to podding plants. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
St John's wort	From flowering to early seed set	All States	500mL	Late spring to early summer.
Sweet briar	Up to 1.5m tall	All States except NT	350mL	Add metsulfuron-methyl at 10g/100L water to obtain more reliable results with the lower rate of Gusto Woody & Noxious Weed Herbicide.
			500mL	Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
Tobacco weed	Actively growing plants	NT, Qld and WA only	300mL	Add a 100% concentrate non-ionic surfactant at 100mL/100L of water for best results.
Wattle (<i>Acacia</i> spp.) (except corkwood wattle)	1 to 3m tall	All States	350mL	
Wild rosemary (<i>Casinia laevis</i>)	Active growth, 0.5 to 1m tall	Qld only	350 to 500mL	Use lower rate on seedlings 0.5m tall. Apply as a thorough foliar spray.
Wild tobacco tree	Spring to autumn up to 2m tall	ACT, NSW and Qld only	350mL	

Table B: Aerial Application
See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY			AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY	
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ ha	CRITICAL COMMENTS
Blackberry	Summer to autumn	NSW, Qld, SA and Vic and WA only	10L	Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before respraying with Gusto Woody & Noxious Weed Herbicide. WARNING: Eucalyptus species up to 4m may be killed if sprayed during this treatment. Mature trees which are 15 to 20m tall may be partially defoliated but are likely to recover.
Gorse		Tas only		
Cockspur thorn Crofton weed Lantana Mistflower	Late autumn	NSW, NT and Qld only (helicopter only)	1.5L plus 7.5L 2,4-D amine (500g/L formulation)	Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150L/ha. Follow up respraying will be required.
Lantana			10L	Helicopter application only.
Rubber vine (Not infected with rust)	When flowering	NT and Qld only (helicopter only)	3L to 5L	Use rates will depend upon the density and height of the rubber vine stand. The higher rates should be used on dense stands, however, complete coverage and penetration may be difficult. Follow up respraying will be required. Any regrowth should be sprayed with a suitable basal bark herbicide.
St John's wort	Flowering to early seed set (Nov-Jan)	NSW only	4L	Helicopter application only. Follow up spraying will be required in the following season.
AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS			AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS	
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ ha	CRITICAL COMMENTS
Parkinsonia	Seedlings, 1-2m tall or 12-24 months old	Qld and NT only (helicopter only)	3L	Add Uptake Spraying Oil at 1L/ha.

Table C: Controlled Droplet Application (CDA)

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY	
WEEDS CONTROLLED	WEED GROWTH STAGE
Blackberry In association with: Docks Ragwort St John's wort Thistles	Summer to autumn

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY		
STATE	RATE/ ha	CRITICAL COMMENTS
All States except NT	Apply undiluted	One application may give satisfactory control but subsequent regrowth and seedlings should be resprayed after hardening off. Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before respraying with Gusto Woody & Noxious Weed Herbicide.

Table D: Low Volume High Concentrate Application Techniques (Gas Powered Gun, Sprinkler Sprayer) See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY	
WEEDS CONTROLLED	WEED GROWTH STAGE
Blackberry	Late spring to autumn
Camphor laurel Cockspur thorn Crofton weed	Less than 1.5m high
<i>Eucalyptus</i> species	Seedlings up to 2m tall
Mistflower	Less than 1.5m high
Sweet briar	1.5m tall, full leaf to ripe fruit
St John's wort	During flowering to early seed set
Wild tobacco tree	Less than 1.5m high

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY		
STATE	RATE/ 10L water	CRITICAL COMMENTS
ACT, NSW, Qld, SA, Tas and WA only	335mL	Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
ACT, NSW and Qld only	500mL	
All States	335mL	Gas Powered Gun only: Apply to actively growing bushes not more than 1.5m tall that have not more than 5 stems from the crown. Gas Powered Gun only: One application should provide control. Minor regrowth and seedlings may be retreated the following summer.
ACT, NSW and Qld only	500mL	
NSW only		
NSW, Tas and Vic only		
ACT, NSW and Qld only		Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.

Table E: Boom Application

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY	
WEEDS CONTROLLED	WEED GROWTH STAGE
Galenia	Fresh growth during spring to summer
Sicklepod	Up to flowering
St John's wort	Flowering to early seed set (Nov-Jan)

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY		
STATE	RATE/ ha	CRITICAL COMMENTS
NSW only	5L	Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
NT and Qld only	3L	DO NOT apply to podding plants. Add a 100% concentrate non-ionic surfactant at 100mL/100L of water.
NSW only	2 to 4L	Use the higher rate on dense infestations and when longer residual control is required. Follow up respraying will be required in the following season.

2 FALLOW SITUATIONS

Table A: Boom Application

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

FALLOW	
WEEDS CONTROLLED	WEED GROWTH STAGE
Blackberry nightshade – Suppression only	10 to 25cm tall, prior to flowering
Camel melon Prickly paddy melon Cucumber melon (<i>Cucumis melo</i>)	From 2 leaf to 50cm diameter
Common sowthistle	From 8 leaf to flowering
Cow vine	From 2 to 5 leaf up to 15cm diameter, prior to flowering
Lucerne (established)	Active growth, 15 to 25cm high, during spring
<i>Polymeria pusilla</i>	2 to 12 leaf up to 20cm diameter, prior to flowering

FALLOW		
STATE	RATE/ ha	CRITICAL COMMENTS
NSW and Qld only	200 to 400mL + 1.2L Glyphosate 450 Herbicide + Adjuvant	FOR USE BY GROUND EQUIPMENT ONLY. Plants must be actively growing. Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5cm diameter for <i>Polymeria pusilla</i>). Refer to Glyphosate 450 label for use of adjuvant. DO NOT plant susceptible crops for up to nine months after application, as specified in General Instructions - Minimum Recropping Periods - Black Cracking Clay Soils, NSW & Qld.
	300 to 500mL + 1.2L Glyphosate 450 Herbicide + Adjuvant	Dry conditions after application will increase the recropping interval.
	200 to 400mL + 1.2L Glyphosate 450 Herbicide + Adjuvant	

Table B: Blanket Wiper Application
See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Bitterbark (<i>Alistonia constricta</i>)	From summer to end of autumn	Qld	1:4 (1 part Gusto Woody & Noxious Weed Herbicide to 4 parts water) 2% solution for spot spraying (eg 100mL Gusto Woody & Noxious Weed Herbicide in 5L water)	For use with blanket wipers only. For best results apply in autumn to tall (> 60cm) plants using two opposite directional passes (up and back). Follow up "missed" plants with a spot spray application. These will be obvious after 6 weeks. Blanket wiper applications can be made in summer when plants are smaller but follow up spraying may be necessary. Do not disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months). Spot spraying "missed" plants: Thoroughly wet all stems and leaves without producing any solution run-off. Avoid any spray reaching the soil surface.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

IN TASMANIA FOR BLACKBERRY

DO NOT treat bushes carrying mature or near mature fruit.

FOR NATIVE VEGETATION

Use of Gusto Woody & Noxious Weed Herbicide on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

WITHHOLDING PERIOD

NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

MINIMUM RECROPPING PERIODS – Black Cracking Clay Soils, NNSW & Qld.

Plant-back periods for crops following the application of Gusto Woody & Noxious Weed Herbicide for rates up to 600mL/ha.				
RATE mL/ha	200	300	400	600
CROP	Months			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba bean	4	4	6	6
Chickpea	4	6	6	6
Lucerne	6	9	9	9

These plant-back periods are based on a normal rainfall pattern. During drought conditions (or when rainfall is less than 100mm for a period of 4 months or greater) the plant-back period may be significantly longer.

Table B: Blanket Wiper Application

Plant-back periods for crops following blanket wiper application.	
CROP	Months
Broadleaf Crops	18
Lucerne	6 This will allow any potential soil residues to dissipate, if any, and allow effective control.

Note: Before using Gusto Woody Weed Herbicide in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product, ie. the product with the longest plant-back period, will determine the time between spraying and planting.

COMPATIBILITY

Follow any regional restrictions, and all directions and restrictions on the label, of any chemical mixed with Gusto Woody & Noxious Weed Herbicide (eg. 2,4-D amine). Gusto Woody & Noxious Weed Herbicide is compatible with the following herbicides: 2,4-D Amine, DP 600, Weed Force Renovate Herbicide, metsulfuron-methyl, Glyphosate 450, Glyphosate 490, Fluroxypyr 200.

Gusto Woody Weed & Noxious Weed Herbicide is compatible with the following adjuvants, as per Directions for Use: Uptake, Pulse Penetrant, Brushwet and non-ionic surfactant (1000g/L)

MIXING

Mix only with water.

Half fill the spray unit with water, and add the required amount of Gusto Woody & Noxious Weed Herbicide. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

APPLICATION

1 WOOD WEED SITUATIONS

Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.

A High Volume Spraying

Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.

Hand Gun

Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500kPa (400 to 500kPa for St John's wort). A spray volume of 3000 to 4000L per infested hectare of 1 to 2 metre high blackberry (30 to 40L/100m²) should be used. Use 2000L of spray mixture/ha of Galenia infestation (ie. 20L/100m² infested area).

Knapsack

Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand gun. A spray volume of 3 to 4L/10m² infested area should be used. A spray volume of 2L/10m² should be used for an area infested with Galenia.

B Aerial Application

Apply in 200L of water/ha using an aircraft to apply 100L per pass on a double overlap pattern using nozzle configurations to produce droplets of 250 to 350 micron diameter. The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets. DO NOT spray when wind exceeds 15km/hr and/or air temperature reaches 35°C.

C Controlled Droplet Application (CDA)

Results similar to high volume spraying can be obtained using Micron Herbi® or similar equipment. Select a nozzle to give a flow rate of 2mL/sec and sweeping action of approximately 1m/sec to ensure a droplet density of 20/cm². Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided by CDA unit.

D Low Volume High Concentrate Application Techniques

Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

Gas Powered Gun: Apply 50mL shots to obtain uniform coverage of 4 to 5m² of surface area of bush. This relates to 20 droplets/cm² of leaf surface.

Sprinkler Sprayer: This technique involves using a micro sprinkler that is connected to a hollow fibreglass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 to 200kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

E Boom Application

Application in a minimum spray volume of 200L/ha for Galenia and St John's wort and 600L of water/ha for Sicklepod. Flat fan nozzles are recommended, using pressure in the range of 200-300kPa. Boom height must be set to ensure double overlap of nozzle patterns.

2 FALLOW SITUATIONS

A Boom Application

Application of Gusto Woody & Noxious Weed Herbicide in a minimum spray volume of 50L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200-300kPa. Boom height must be set to ensure double overlap of nozzle patterns.

B Blanket Wiper Application

Blanket needs to be made from durable and wettable material with a rigid backing. Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (rewetting). Ideally, a scraper bar should be mounted in front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping the stems. This scraper may be mounted at the front of the vehicle. Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10-15kph are ideal for blanket wiping application.

CLEANING SPRAY EQUIPMENT

Rinsing

After using Gusto Woody & Noxious Weed Herbicide, empty spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles. After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

Decontamination

Before spraying cotton and other sensitive crops with equipment that has been used to apply Gusto Woody & Noxious Weed Herbicide, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section. Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (eg. liquid SURF, OMO, OMOMATIC, DRIVE at 500mL/100L of water or the powder equivalent at 500g/100L of water) and circulate throughout the system for at least 15 minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

RESISTANT WEEDS WARNING

GROUP 4 HERBICIDE

Weed Force Gusto Woody & Noxious Weed Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group 4 herbicide.

Some naturally occurring weed biotypes resistant to the product and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group 4 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Weed Force Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Weed Force representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Crops susceptible to Gusto Woody & Noxious Weed Herbicide include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and Pinus spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines. Gusto Woody Weed & Noxious Weed Herbicide

is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment. Picloram, one of the active constituents in this product, can remain in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation. DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops. DO NOT apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants. DO NOT apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants. DO NOT move soil, which may have been treated to areas where desirable plants are to be grown.

PROTECTION OF LIVESTOCK

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.

Many plants remain poisonous after death, and stock should not be allowed access to it, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well-ventilated area, as cool as possible out of direct sunlight. DO NOT store or expose product to wet conditions.

Keep away from contact with fertilisers, fungicides and seeds.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb as above, the wash liquid for disposal.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (PHONE AUSTRALIA: 13 1126).
If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

The SDS contains important information relating to health and safety. This product should be used in conjunction with information on the SDS and users are advised to read the SDS before using this product. The most up to date SDS can be obtained from www.weedforce.com.au

CONDITIONS OF SALE

To the extent permitted by law, all conditions and warranties and statutory or other rights or action which buyer or any other user may have against Weed Force Pty Ltd ("Weed Force") or Seller are hereby excluded. Weed Force hereby gives Notice to Buyer and other users that it will not accept responsibility for any direct, indirect or consequential loss however arising (including negligence) from the sale, supply, use or application of this product or reliance on product information or advice provided by Weed Force or on its behalf. No warranty expressed or implied is given by Weed Force regarding its suitability or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not. Weed Force's liability shall, in all circumstances, be limited to replacement of the product or a refund of the purchase price paid therefor.



WEEDFORCE Pty Ltd ACN 602 207 152 
2072 Logan Rd Mt Gravatt QLD 4122
P: 0488 997 923 www.weedforce.com.au

APVMA Approval No:
83353/111562

® Trademark of Weed Force Pty Ltd
© Weed Force Pty Ltd 2017